

Which Higgs-Yukawa systems can possess non-trivial fixed points

Sergei V. Zenkin^a

^aInstitute for Nuclear Research of the Russian Academy of Sciences, 60th October Anniversary
Prospect 7a, 117312 Moscow, Russia

We argue that non-trivial fixed points bordering on the paramagnetic and ferromagnetic phases are most likely to exist in the Higgs-Yukawa systems that have a connected domain with the paramagnetic phase and no ferrimagnetic phase. We find three examples of such systems; among them is the U(1) system with naive fermions.

1. In this talk we want to emphasize the following three points:

(i) that the phase structure of the Higgs-Yukawa systems depend crucially both on their symmetry group and on the form of the lattice fermion action, even though the coupling of the fermions to the Higgs fields is the same;

(ii) that the most likely candidates for the systems with non-trivial fixed points bordering on the paramagnetic (PM) and ferromagnetic (FM) phases are the systems that have a connected domain with the PM phase and no ferrimagnetic (FI) phase;

(iii) that one of such systems is the U(1) system with naive fermions.

We investigate the phase diagrams of the Z_2 , U(1) and SU(2) Higgs-Yukawa systems with radially frozen Higgs fields for three types of chirally invariant lattice fermion actions. Our method is based on the variational mean field approximation, where contribution of the fermion determinant is calculated for weak and strong Yukawa coupling regimes in a certain ladder approximation [1,2].

2. The action of the system

$$A = - 2\kappa \sum_{n,\mu,a} \phi_n^a \phi_{n+\hat{\mu}}^a + \sum_{m,n} \bar{\psi}_m (D_{mn} + y \tilde{\Phi}_m \delta_{mn}) \psi_n \quad (1)$$

has two parameters: the scalar hopping parameter κ and the Yukawa coupling y ; ϕ^a are real components of the group-valued field Φ , $\tilde{\Phi} = P_L \Phi^\dagger + P_R \Phi$. Given group, the system is deter-

mined by the form of the lattice Dirac operator D . At $\kappa > 0$ it corresponds to certain continuum Higgs-Yukawa model; at non-positive κ such a correspondence is destroyed.

We shall require the system to be chirally invariant and operator D to have the form:

$$\begin{aligned} D_{mn} &= -D_{nm}, \\ D_{mn} &= \int_p e^{ip(m-n)} i \gamma_\mu L_\mu(p), \end{aligned} \quad (2)$$

with real $L_\mu(p)$; $\int_p \equiv \int d^4p/(2\pi)^4$, $p_\mu \in (-\pi, \pi)$.

We consider three examples of the lattice fermions satisfying these conditions: naive fermions with

$$L_\mu(p) = \sin p_\mu; \quad (3)$$

non-local SLAC fermions with

$$L_\mu(p) = p_\mu; \quad (4)$$

and fermions with

$$L_\mu(p) = \sin p_\mu \left[1 + \frac{(2 \sum_\nu \sin^2 p_\nu / 2)^2}{\sum_\nu \sin^2 p_\nu} \right], \quad (5)$$

whose action although is non-local, is originated from the local mirror fermion action after integrating out the mirror fermions (see [3] and also [2]).

In all the cases the fermions couple to the Higgs fields in the same way.

3. The method [1,2] yields closed analytical expressions for the critical lines $\kappa(y)$ between the FM and PM phases and between the anti-ferromagnetic (AM) and PM phases in the weak

(W) and strong (S) coupling regimes: $\kappa^W(y)$ and $\kappa^S(y)$, respectively. Their explicit form for the Z_2 , $U(1)$ and $SU(2)$ systems can be found in [2]. The important fact is that for given group, the form of these line is mainly determined by four constants: $G^W(0)$, $G^W(\pi)$, $G^S(0)$ and $G^S(\pi)$, which are the values of the functions

$$\begin{aligned} G^W(q) &= \int_p \frac{L(p)L(p+q)}{L^2(p)L^2(p+q)}, \\ G^S(q) &= \int_p L(p)L(p+q) \end{aligned} \quad (6)$$

at $q = 0$ and $q = \pi$. Moreover, in the cases of Z_2 and $SU(2)$ these constants determine singular points of the expressions for $\kappa^W(y)$ and $\kappa^S(y)$, thereby determining the domains of the weak and strong coupling regimes in these cases. These domains may be separated or they may overlap. Both possibilities are realized in our examples.

As a result we get three types of the phase diagrams [2]:

(I) If these domains are separated and FM-PM critical lines do not intersect AM-PM lines, we have the diagram with two domains with PM phase separated by the funnel with the FM phase at the intermediate values of y . This is the case of Z_2 systems with the naive and mirror fermions.

(II) If the FM-PM lines intersect AM-PM lines before possible intersection of the lines $\kappa^W(y)$ and $\kappa^S(y)$, FI phase arises and PM phase occupies two disconnected domains, too. This is the case for the $U(1)$ systems with more than one SLAC fermions or with mirror fermions and for all the $SU(2)$ systems.

(III) Finally, if the domains of weak and strong coupling regimes overlap, or they are not determined, like in the $U(1)$ case, and the FI phase does not arise, we have the diagram shown in Fig. 1. Domain with the PM phase is now connected. This is the case for the $U(1)$ systems with naive or one SLAC fermions and for Z_2 system with SLAC fermions.

In all the cases fermion correlators have different properties in the weak and strong coupling regimes, at least in the FM and PM phases, thereby distinguishing FMW and FMS, and PMW and PMS phases [5] (see also [2]). In particular, in PMW phase the fermions are mass-




Figure 1. Phase diagram of the type (III) (solid lines); fermion condensate along the FM-PM critical line (dashed lines).

less, while in PMS phase they are decoupled from the low lying spectrum.

4. We now concentrate on the neighbourhood of the FM-PM critical lines, since it is this domain is the most interesting for application to the elementary particle physics. Consider the fermion condensate along these lines. In our approximation this quantity has the form

$$\langle \bar{\psi} \psi \rangle = -2^{D/2} C(y) \langle \phi \rangle + O(\langle \phi \rangle^2), \quad (7)$$

where $\langle \phi \rangle$ is v.e.v. of the Higgs field, and for given group function $C(y)$ is determined in the weak and strong coupling regimes by the constants $G^W(0)$ and $G^S(0)$, respectively, [2].

The important fact is that everywhere along the FM-PM critical lines the condensate turns out to be continuous smooth function of y , except for the only point A in the phase diagram of the type (III) (see Fig. 1). In this figure dashed lines show the function $C(y)$ for the $U(1)$ system with naive fermions. The condensate can be defined as the first order derivative of free energy of the system. Therefore, its discontinuity in the point A is an indication to a first order phase transition separating FMW and FMS (and perhaps PMW and PMS) phases. In Fig. 1 it is shown by the crossed line. Since the condensate is an order parameter of the system, too, the point A looks



Figure 2. Possible variant of the phase diagram of the type (III).

like tricritical point at which the first order phase transition becomes a second order one.

Thus, we come to the scenario discussed in ref. [4]. In view of the results of ref. [7] that have given no evidence of non-trivial behaviour of the SU(2) system in the point of forming the FI phase, the point A is the only candidate for non-trivial fixed point bordering on the FM and PM phases.

5. So, we have found that only three of the considered systems possess such points. Note, that all these points lie at negative κ , where the systems have no their continuum counterparts and are not reflection positive. Therefore, the necessary condition for the systems to be of interest for continuum physics is their physical positivity at $\kappa < 0$. However, if this condition is satisfied and some of these points turn out to be indeed non-trivial, the problem of universality of the Higgs-Yukawa systems will arise. In any case, it would be interesting to find the systems where such points are located at $\kappa > 0$.

Among these three systems is the U(1) system with the naive fermions. The coordinates of the point A in this case is: $y^A \approx 1.34$, $\kappa^A \approx -0.43$ (for two fermions). The existence of such a point in this system, however, disagrees with Monte Carlo (MC) results of ref. [6], where FI phase has

been observed. We believe that this disagreement is due to the finite lattice effects in the MC calculations. Our results for SU(2) system with naive fermions and for U(1) and SU(2) systems with mirror fermions are in agreement with MC results [5,3]. In these cases the reasons for appearance of the FI phase are quite clear. Therefore we see no reasons why our method should fail in this case.

The only change in the diagrams of the type (III) which we cannot exclude is that the FM-PM critical line (and, perhaps, AM-PM line, too) actually has a discontinuity at some value of y not far from y^A . In this case the point A will be shifted and split into two points, for example, as it is shown in Fig. 2. Investigation of this question, however, requires different methods.

ACKNOWLEDGMENTS

I am grateful to the International Science Foundation and to Organizing Committee of the "Lattice '94" for financial support, and to Ph. Boucaud and to M. Stephanov for interesting discussions.

REFERENCES

1. S. V. Zenkin, Mod. Phys. Lett. A9 (1994) 983.
2. S. Tominaga, S. V. Zenkin, Phys. Rev. D50 (1994) 3387.
3. L. Lin et al., Phys. Lett. B264 (1991) 407; C. Frick et al., Nucl. Phys. B397 (1993) 431.
4. A. Hasenfratz and T. Neuhaus, Phys. Lett. B220 (1989) 435.
5. W. Bock et al., Nucl. Phys. B344 (1990) 207.
6. A. Hasenfratz et al., Phys. Lett. B236 (1990) 339.
7. W. Bock et al., Nucl. Phys. B371 (1992) 683.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-lat/9411069v1>